

12 LEVEL


AD-E430146

TECHNICAL REPORT ARBRL-TR-02106

AN ANALYSIS OF THE PRESSURE DEPENDENCE  
OF NITRATE ESTER THERMAL DECOMPOSITION

George F. Adams

September 1978



AD A063513

DDC FILE COPY



US ARMY ARMAMENT RESEARCH AND DEVELOPMENT COMMAND  
BALLISTIC RESEARCH LABORATORY  
ABERDEEN PROVING GROUND, MARYLAND

Approved for public release; distribution unlimited.

78 11 20 037

Destroy this report when it is no longer needed.  
Do not return it to the originator.


Secondary distribution of this report by originating  
or sponsoring activity is prohibited.

Additional copies of this report may be obtained  
from the National Technical Information Service,  
U.S. Department of Commerce, Springfield, Virginia  
22161.

The findings in this report are not to be construed as  
an official Department of the Army position, unless  
so designated by other authorized documents.

*The use of trade names or manufacturers' names in this report  
does not constitute endorsement of any commercial product.*

# DISCLAIMER NOTICE



THIS DOCUMENT IS BEST  
QUALITY AVAILABLE. THE COPY  
FURNISHED TO DTIC CONTAINED  
A SIGNIFICANT NUMBER OF  
PAGES WHICH DO NOT  
REPRODUCE LEGIBLY.

REPRODUCED FROM  
BEST AVAILABLE COPY

THIS DOCUMENT CONTAINED  
BLANK PAGES THAT HAVE  
BEEN DELETED

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report ARBRL-TR-02106	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER Technical Rept.
4. TITLE (and Subtitle) An Analysis of the Pressure Dependence of Nitrate Ester Thermal Decomposition,	5. TYPE OF REPORT & PERIOD COVERED	
7. AUTHOR(s) George F. Adams	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Ballistic Research Laboratory ATTN: DRDAR-BLP Aberdeen Proving Ground, MD 21005	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1L161102AH60	
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Armament Research & Development Command US Army Ballistic Research Laboratory ATTN: DRDAR-BL, APG, MD 21005	12. REPORT DATE SEPTEMBER 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) OSBIA	13. NUMBER OF PAGES 34	
16. DISTRIBUTION STATEMENT (for this report) Approved for public release; distribution unlimited.	15. SECURITY CLASS. (of this report) Unclassified	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) N-propyl nitrate, thermal decomposition, unimolecular reaction, nitrate ester decomposition, solid propellant, combustion.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) (hmn) An analytical study of the effect of pressure on the rate constant for the unimolecular decomposition of n-propyl nitrate is given. The Rice-Ramsperger-Kassel-Marcus theory is reviewed, and the present application is discussed in detail. Limitations on application of this model are also discussed. It is concluded that the initial decomposition reaction of nitrate esters cannot be the rate determining step for the observed variation of burning rate with pressure.		

DDC  
RECEIVED  
JAN 22 1979  
B

# TABLE OF CONTENTS

	Page
LIST OF SYMBOLS . . . . .	5
I. INTRODUCTION. . . . .	7
II. UNIMOLECULAR REACTION RATE THEORY . . . . .	7
III. RRKM PARAMETERS FOR N-PROPYL NITRATE. . . . .	11
IV. RESULTS OF THE RRKM INTEGRATION AND COMPARISON WITH BURNING-RATE DATA . . . . .	15
REFERENCES. . . . .	21
APPENDIX A. COMMENTS ON DETAILS OF THE RRKM CALCULATION. . . . .	23
APPENDIX B. DATA FORMAT FOR RRKM . . . . .	25
REFERENCES. . . . .	28
DISTRIBUTION LIST . . . . .	29

ACCESSION for	
NTIS	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
DDC	<input type="checkbox"/> Yes <input type="checkbox"/> No
UNANNOUNCED	<input type="checkbox"/> Yes <input type="checkbox"/> No
JUSTIFICATION	
BY	
DISTRIBUTION	CODES
Dist.	RECEIVED
A	

# LIST OF SYMBOLS

$h, k, N_A, R$	Planck, Boltzman, Avogadro and gas constants.
$p, T$	pressure and absolute temperature
$Q$	partition function
$A^*, A^\ddagger$	energized molecule and activated complex
$E^* (\equiv E_{VR}^*)$	total non-fixed energy in the active degrees of freedom of a given energized molecule.
$E_v^*, E_r^*$	vibrational and rotational parts of $E^*$
$E^\ddagger$	total non-fixed energy in the active degrees of freedom of a given activated complex
$E_v^\ddagger, E_r^\ddagger$	vibrational and rotational contributions to $E^\ddagger$
$k_1$	rate-constant for energizing of reactant
$k_a$	rate-constant for conversion of energized molecule to products, Lindeman theory.
$k_a(E^*)$	same rate-constant with energy-dependence, as in RRKM theory
$k_2$	rate constant for de-energizing of energized molecule
$E_0$	critical energy for reaction
$k_{uni}$	first order rate constant, $-(1/[A])d[A]/dt$
$L^\ddagger$	statistical factor or reaction path degeneracy
$Q_1, Q_1^\ddagger$	partition functions for adiabatic rotations in $A$ and $A^\ddagger$
$Q_2$	partition functions for active vibrational degrees of freedom in $A$
$\sum P(E_{VR}^*), W(E_{VR})$	number of quantum states of $A^\ddagger$ at all energies up to and including $E^\ddagger$ .
$N^*(E^*)$	density of quantum states of $A^*$ at energy $E^*$ .
$\lambda$	collisional energy transfer efficiency
$Z$	second-order collision rate constant
$\Delta S^\ddagger$	entropy of activation.

## I. INTRODUCTION

Several recent models have explained the effect of pressure on ballistic properties as due to an assumed change in the microscopic processes that determine the extensive properties (1,2). Each model assumes that an initial unimolecular decomposition reaction is the rate-determining step over the entire pressure regime peculiar to the macroscopic process. Both models assume that external pressure causes a major change in the rate of this decomposition reaction, thereby controlling the ballistic phenomena. This report describes an analytical study of the effect of pressure on the thermal decomposition of a nitrate ester, n-propyl nitrate. The results of the study demonstrate that the pressure-dependence of this reaction rate cannot explain the observed burning rate behavior of nitrate esters.

The second section of this report describes the chemical reaction being investigated and the methods used to perform the analytical study. This research involves the application of the Rice-Ramsperger-Kassel-Marcus (RRKM) theory to predict the rate constant for the decomposition reaction as a function of pressure. The third section describes the parameters used to describe the reacting system, and provides details of the kinetics calculations. A final section provides a comparison of the RRKM results and the observed burning-rate behavior, and contains comments on the validity of the assumptions that underly the RRKM theory. Several appendices provide information about portions of the calculation that deserve detailed explanation.


## II. UNIMOLECULAR REACTION RATE THEORY

N-propyl nitrate was chosen for study because of good experimental data on the thermal decomposition reaction. The initial decomposition reaction is identified as the scission of the nitro-alkoxy bond to produce two free radicals,  $\text{NO}_2$  and alkoxy. This unimolecular decomposition has been studied for n-propyl nitrate using the very low pressure pyrolysis (VLPP) method (3). An RRKM analysis of the temperature dependence of the high pressure rate constant has been performed but no analysis of the pressure-dependence of the unimolecular reaction rate was made (3).

1. M.M. Ibricu and F.A. Williams, "Mechanisms for the Steady Deflagration of Double-Base Propellants", 12th JANNAF Combustion Meeting, CPIA Publication Number 273, 1976.
2. D.J. Pastine, M.J. Kamlett and S.J. Jacobs, "Volume and Pressure Dependence of Some Kinetic Processes in Explosives", Naval Surface Weapons Center/WOL/TR75-149, 1975.
3. G.D. Mendenhall, D.M. Golden, and S.W. Benson, "The Very-Low-Pressure Pyrolysis (VLPP) of n-Propyl Nitrate, tert-Butyl Nitrate, and Methyl Nitrite. Rate Constants for Some Alkoxy Radical Reactions", Int. J. Chem. Kinetics, 7, 725 (1975).

The present research consists of an RRKM analysis of the pressure dependence of the nitrate ester bond-scission reaction. The following paragraphs outline the development of RRKM theory. More complete discussions can be found in books by Forst (4), and Robinson and Holbrook (5).

A unimolecular decomposition reaction involves a single molecule that undergoes transformation to a reactive complex by some energization process. The formation of the complex is a function of intermolecular energy transfer, often via collision, and intramolecular energy transfer after the collision occurs. Thus, the process is envisioned as



Most modern theories that describe unimolecular decomposition reactions are based on the Lindeman mechanism (6) which assumes that collisions between two reactant molecules will produce some energized molecules with enough energy to react. If conversion of reactant molecules to energized molecules is slow compared to the rate at which collisional de-energization occurs, an equilibrium population of energized molecules will exist with concentration proportional to the concentration of the reactant molecules. Thus the rate constant will be first-order with respect to reactant concentration. At low pressure, however, collisions cannot maintain an equilibrium concentration of energized intermediates, so the rate constant will depend on the rate of energization which is proportional to the square of the concentration of reactant molecules. Lindeman's mechanism predicts that at low pressure the rate constant increases with increasing pressure, but at high pressure the rate constant reaches a limit and remains constant. Although these predictions are qualitatively correct, numerical studies based on Lindeman's mechanism yield poor agreement between theory and experiment.



Hinshelwood extended Lindeman's theory and laid the basis for the development of RRKM theory by suggesting that one needs to consider the internal degrees of freedom when determining whether a molecule possesses sufficient energy to react (7). The chance for a molecule to have enough energy to react increases with the number of degrees of freedom of the reactant molecule, primarily because of the contribution of the vibrational degrees of freedom.

4. W. Forst, Theory of Unimolecular Reactions, Academic Press, New York, (1973).
5. P.J. Robinson and K.A. Holbrook, Unimolecular Reactions, Wiley-Interscience, New York (1972).
6. F.A. Lindeman, title not available, Trans. Faraday Soc., 17, 598 (1922).
7. C.N. Hinshelwood, "The Theory of Unimolecular Reactions", Proc. Roy.Soc. (A), 113, 230 (1927).



The final important assumption, suggested simultaneously by Rice and Ramsperger (8,9) and Kassel (10), is that the rate of decomposition of the energized molecule is a function of the energy of the species. The species will not decompose unless it possesses a minimum internal energy, called the critical energy  $E_0$ , and will react at a rate that increases as a function of the energy it contains in excess of  $E_0$ . The corollary of this is that if energy is the necessary and sufficient condition for reaction, it is implied that energy is exchanged among the internal degrees of freedom. That is, there is intramolecular energy transfer. The developers of RRKM theory ignore the dynamics of such transfer, assume that the transfer is a random process, and go on to develop a statistical theory couched in the language of transition-state theory (11).

The reaction scheme for the RRKM model is



The notation  $E^* \rightarrow E^* + \delta E^*$  means that the energized molecule has energy in the range  $E^*$  to  $E^* + \delta E^*$ . The second equation represents the major feature of the RRKM theory wherein Absolute Rate Theory is applied to the calculation of  $k_a(E^*)$ . Thus we distinguish between an energized molecule  $A^*$  and the activated complex  $A^\ddagger$ .

8. O.K. Rice and H.C. Ramsperger, "Theories of Unimolecular Reactions at Low Pressures", J. Amer. Chem. Soc., 49, 1617 (1927).
9. O.K. Rice and H.C. Ramsperger, "Theories of Unimolecular Reactions at Low Pressures - II", J. Amer. Chem. Soc., 50, 617 (1928).
10. L.S. Kassel, Kinetics of Homogeneous Gas Reactions, Chemical Catalog Co., New York, 1932.
11. R.A. Marcus, "Unimolecular Dissociations and Free Radical Recombination Reactions", J. Chem. Phys., 20, 359 (1952).

The energized molecule  $A^*$  is a molecule that contains in its active degrees of freedom a non-fixed energy<sup>#</sup>  $E^*$  greater than the critical energy  $E_0$ . This non-fixed energy may consist of both vibrational and rotational energy.

The activated complex  $A^\ddagger$  is a species that is recognizable as intermediate between reactants and products, and is characterized by a configuration that corresponds to a position atop a potential barrier. The total non-fixed energy of the activated complex is  $E^\ddagger$  which is equal to  $E^*$  less the critical energy  $E_0$ .

The RRKM theory provides the following mathematical expression for the overall unimolecular rate-constant,  $k_{uni}$ ,

$$k_{uni} = \int_{E^* = E_0}^{\infty} \frac{k_a(E^*) dk_1(E^* \rightarrow E^* + \delta E^*)/k_2}{1 + k_a(E^*)/k_2[M]} \quad (4)$$

Using statistical mechanics, expressions for  $\delta k_1(E^* \rightarrow E^* + \delta E^*)/k_2$  and  $k_a(E^*)$  are obtained leading to

$$k_{uni} = \frac{L^\ddagger Q_1^\ddagger}{h Q_1 Q_2} \int_{E^* = E_0}^{\infty} \frac{\{\sum P(E_{VR}^\ddagger)\} \exp(-E^*/kT) dE^*}{1 + k_a(E^*)/k_a[M]} \quad (5)$$

with

$$k_a(E^*) = \frac{L^\ddagger Q_1^\ddagger}{h Q_1 N^*(E^*)} \sum_{E_{VR}^* = 0}^{E^*} P(E_{VR}^*). \quad (6)$$

where

- $Q_1^\ddagger$  rotational partition function for the activated complex
- $Q_1$  rotational partition function for the reactant
- $Q_2$  vibrational partition function for the reactant
- $h$  Planck's constant
- $\sum P(E_{VR}^\ddagger)$  number of quantum states of  $A^\ddagger$  at all energies up to  $E^\ddagger$
- $N^*(E^*)$  density of quantum states of  $A^*$  at energy  $E^*$
- $L^\ddagger$  statistical factor or reaction path degeneracy.

<sup>#</sup>Non-fixed energy is internal energy available for redistribution.

In practice the product  $k_2[M]$  is replaced by the collisional deactivation rate,

$$k_2[M] = \lambda Z p \quad (7)$$

where

$\lambda$ , collisional efficiency  
 $Z$ , collision number  
 $p$ , pressure.

The RRKM integral is rewritten as a sum, resulting in the following working equation for the evaluation of  $k_{uni}$ .

$$k_{uni} = \frac{L^{\dagger} Q_1^{\dagger}}{h Q_1 Q_2} \exp(-E_o/RT) \Delta E^{\dagger} \sum_{L=1}^{i_{max}} \left[ \frac{\{ \sum P(E_{VR}^{\dagger}) \} \exp(-E^{\dagger}/RT)}{1 + k_a(E^{\dagger})/\lambda Z p} \right]_i \quad (8)$$

where the energy relations,

$$E^{\dagger} = E_o + E^{\dagger}, \quad E^{\dagger} = (i + \frac{1}{2}) \Delta E_i \quad (9)$$

have been used. The rate constant will have units  $\text{sec}^{-1}$  if the energies are expressed in  $\text{kcal mol}^{-1}$  and the constants  $h$  and  $R$  have the values

$$h = 9.537 \times 10^{-14} \text{ kcal mol}^{-1} \text{ sec} \quad (10)$$

$$R = .0019872 \text{ kcal mol}^{-1} \text{ K}^{-1}. \quad (11)$$

The expressions given in equations 6 and 8 are the expressions used in our study of the decomposition of n-propyl nitrate. In the following section the methods used to evaluate variables in these expressions are reviewed and the model used for the n-propyl nitrate calculations is specified.

### III. RRKM PARAMETERS FOR N-PROPYL NITRATE

There are a number of variables that must be determined before a prediction of  $k_{uni}$  can be made. The computation of these variables requires input information about the structure and properties of the reactant molecule and the activated complex. In general, information about the reactant is available, whereas information about the complex is unknown. Important data that are required include:

1. Vibrational frequencies
2. Moments of inertia
3. Collision number
4. Critical energy,  $E_0$ .

Vibrational frequencies for the reactant are needed for calculation of the vibrational partition function,  $Q_2$ , and the density of quantum states,  $N^*(E^*)$ . Vibrational frequencies for the activated complex are needed in order to calculate the sum of quantum states of  $A^*$ .

The vibrational frequencies for N-propyl nitrate are known, but those for a transition state are neither known nor measurable. Often, vibrational frequencies for the transition state can be assigned by transfer of data from molecules with analogous structures. In the case of bond stretching frequencies, bond force constants are predicted on the basis of bond-order correlations (12). However, there is no well-established method for estimating the force constants for bending and torsional modes.

Benson and O'Neal have tabulated frequencies for stretching and bending modes and used these values for predicting the high-pressure A-factors of unimolecular reactions (13,14).

Any attempt to deduce the properties of the activated complex leaves some of the variables undetermined. Generally, these variables are used as adjustable parameters to make the model predict the correct value for the experimental high-pressure frequency factor. Mendenhall, Golden and Benson (3) have made no effort to deduce the properties of the complex since it is known that the calculated fall-off behavior is insensitive to the model used provided that it predicts the correct value of  $A_\infty$ . The high-pressure rate constant,  $k_\infty$ , is given in RRKM theory by the expression,

$$k_\infty(\text{RRKM}) = L^\ddagger \frac{kT}{h} \frac{Q_1^\ddagger Q_2^\ddagger}{Q_1 Q_2} \exp(-E_0/kT) \quad (12)$$

12. H.S. Johnston, Gas-Phase Reaction Rate Theory, Ronald, New York, 1966.
13. S.W. Benson, Thermochemical Kinetics, Wiley, New York, 1968.
14. H.E. O'Neal and S.W. Benson, "The Biradical Mechanism in Small Ring Compounds", J. Phys. Chem., 72, 1866, 1968.

This implies that the logarithm of the high-pressure frequency factor is given by

$$\ln A_{\infty}^{\ddagger}(\text{RRKM}) = \ln (L^{\ddagger} e kT/h) + d \left( T \ln \frac{Q_1^{\ddagger} Q_2^{\ddagger}}{Q_1 Q_2} \right) / dT \quad (13)$$

$$= \ln(L^{\ddagger} e kT/h) + \frac{\Delta S^{\ddagger}}{R} \quad (14)$$

A convenient numerical value of this expression is

$$\log_{10} A_{\infty}^{\ddagger} = 10.753 + \log_{10}(L^{\ddagger} T) + \Delta S^{\ddagger} / 4.756 \quad (15)$$

In practice, the value of  $\Delta S^{\ddagger}$  is adjusted so that the high-pressure frequency factor agrees with the experimental value. The change in entropy is divided into vibrational and rotational contributions;

$$\Delta S_V^{\ddagger} = S_V(A^{\ddagger}) - S_V(A) \quad (16)$$

$$\Delta S_R^{\ddagger} = S_R(A^{\ddagger}) - S_R(A) \quad (17)$$

To solve for these contributions it is assumed that the vibrations are quantum harmonic oscillators and the rotations are classical. The statistical mechanical expression for the entropy is

$$S = R \ln Q + RT \frac{d \ln Q}{dT} \quad (18)$$

The appropriate expressions to calculate the vibrational and rotational entropy are

$$Q_{V_i} = \left[ 1 - \exp(-h \nu_i / kT) \right]^{-1} \quad (19)$$

$$T \frac{d \ln Q_{V_i}}{dT} = \frac{(h \nu_i / kT)}{[\exp(h \nu_i / kT) - 1]} \quad (20)$$

$$Q_R = \pi^{1/2} (8 \pi^2 kT/h)^{3/2} (I_a I_b I_c)^{1/2} \quad (21)$$

$$T \frac{d \ln Q_R}{dT} = 3/2 \quad (22)$$

Given these expressions, the use of the RRKM method to study unimolecular decomposition kinetics is straightforward. The molecular parameters chosen by Mendenhall, et al, are given in Table I with one exception. In this study, the RO-NO<sub>2</sub> distance was reduced by .06 Å in both the reactant and the complex. This change has almost no effect on the overall calculation. The parameters listed yield a value for the logarithm of the high-pressure A-factor of 16.5. This relatively high-value is expected because the nitro-moiety becomes a free-rotor in the transition state.

In order to complete the RRKM integration as a function of pressure, one must specify the collision number, Z, and the collisional efficiency, λ. The experimental apparatus used by Mendenhall, et al, (3), is claimed to have a collisional efficiency near one. The collision number, Z, is given by the expression

$$Z = (\sigma_d^2 N_A / R) (8\pi N_A k / \mu)^{1/2} (1/T)^{1/2}, \quad (23)$$

with Z in units Torr<sup>-1</sup> sec<sup>-1</sup> when:

$\sigma_d$  = collision diameter (cm)

$\mu$  = reduced molar mass (g mol<sup>-1</sup>)

T = temperature (K)

$N_A$  = 6.0225 \* 10<sup>23</sup> mol<sup>-1</sup>

R = 6.2362 \* 10<sup>4</sup> cm<sup>3</sup> Torr K<sup>-1</sup> mol<sup>-1</sup>

k = 1.3805 \* 10<sup>-16</sup> erg K<sup>-1</sup>.

The temperature used in this study was 600°K, a temperature assumed to be the surface temperature of a burning double-base propellant. The collision diameter,  $\sigma_d$ , was chosen to be 10<sup>-7</sup> cm, slightly greater than the distance between the most separated atoms in the equilibrium structure of n-propyl nitrate.

#### IV. RESULTS OF THE RRKM INTEGRATION AND COMPARISON WITH BURNING-RATE DATA

The data given in Table I have been used to estimate values for  $k_a(E^*)$  and  $\sum P(E_{VR}^+)$  using the Hase-Bunker computer code (15). These results and the collision number have been used to evaluate  $k_{uni}$  as a function of pressure - using a desk-top computer (Tektronix 4051) to calculate the rate constant. Table II lists the value for  $k_{uni}$  as a function of pressure, and the results are shown in Figure 1. It is apparent from the figure that the unimolecular rate constant is a constant for pressures above  $\sim 400$  Torr, or slightly more than one-half atmosphere. Thus, this research predicts that the rate constant for the initial scission reaction of this medium-size nitrate ester is independent of external pressure for pressures above one atmosphere, for temperatures comparable to those at the surface of a burning propellant. What does this imply in regard to the burning of nitro-cellulose?

Nitrocellulose, a polymeric material, possesses an indeterminately large number of vibrational frequencies. It is a characteristic of unimolecular reactions that as molecules become large, the effect of pressure on the rate constant disappears; that is, the fall-off region moves to lower pressure. Consequently, we expect that the effect of pressure on  $k_{uni}$  for nitrocellulose decomposition is negligible. Note, however, that we mean only the initial scission of the nitrate ester; our results cannot be extended to the decomposition reactions of secondary radicals formed in the overall chemical kinetic scheme of nitrocellulose combustion. The next report in this series will demonstrate the crucial role played by pressure-variation for the decomposition of small radicals.

Given this demonstration of the virtual pressure-independence of the nitrate-ester bond-scission reaction, it can be concluded that the pressure-dependence of propellant burning-rate cannot be explained on the basis of this reaction. This conclusion, however, must be tempered by a consideration of the assumptions made in developing RRKM theory, assumptions that limit the applicability of the theory. There are two assumptions made in the statistical development of RRKM theory.

The first assumption is a corollary of the assumption that energy is the necessary and sufficient condition for reactions, it is that the internal energy of the energized molecule is randomly distributed. If the energy is not randomly distributed, then the high

15. W.L. Hase and D. Bunker, RRKM, Program No. 234, Quantum Chemistry Program Exchange, University of Indiana, Bloomington, Indiana.




Figure 1. Plot of the unimolecular rate constant,  $k_{uni}$ , versus log of pressure for N-propyl nitrate.



Table I. Molecular Parameters for N-Propyl Nitrate.

Vibrational frequencies, $\text{cm}^{-1}$ , with degeneracy in parentheses	Molecule	Complex
	3000 (7)	3000 (7)
	1600	1600
	1450 (4)	1450 (4)
	1350 (4)	1350 (4)
	1200	1150 (5)
	1000	1000 (2)
	1150 (4)	720 (2)
	750 (3)	
	350 (4)	
	250 (3)	125 (6)
	100	50 (3)
	50 (2)	
RO-NO <sub>2</sub> , bond length, Å°	1.407	2.407
Moments of Inertia, amu Å° <sup>2</sup>		
I <sub>a</sub>	58.1	68.5
I <sub>b</sub>	412.3	542.8
I <sub>c</sub>	460.8	601.7
I <sub>red</sub>		39.1

Table II. Rate Constant versus Pressure Data for 600°K Decomposition of N-Propyl Nitrate.

Pressure (torr)	k <sub>uni</sub>
0.5	24.858
1.	30.559
10.	44.088
20.	45.902
100.	47.615
200.	47.849
400.	47.969
1000.	48.04
10000.	48.08
100000.	48.09

pressure rate constant will not be constant, but will show an increase over a finite range of pressure above the limit for the decomposition of molecules with randomly distributed internal energy. The effect of non-random distribution of energy has been discussed by Oref (16) and commented on by Adams and Ward (17). It is unlikely that such an effect will be observed in molecules such as n-propyl nitrate or nitrocellulose, since the temperature remains relatively low during the decomposition process.

The second assumption made in developing RRKM theory is that a single vibrational potential defines the reaction hypersurface. This means, in dynamic terms, that the entire reaction proceeds along a pathway corresponding to a single potential energy surface. The problem can be understood with the aid of Figure 3, a schematic that shows the electronic energy for the first three electronic states of a nitrate ester plotted as a function of the alkoxy-nitro bond length. The curve labeled  $W_1$  corresponds to the energy of the electronic ground state, while  $W_2$  and  $W_3$  label the first two triplet excited states. (This figure is based on assignments first discussed by L.E. Harris (18)). Curves  $W_1$  and  $W_2$  are everywhere separated by an energy difference, so the state represented by  $W_2$  cannot influence the decomposition of a nitrate ester in state  $W_1$ . (It is expected that all the nitrate ester molecules are initially in the state represented by  $W_1$ ). If only states  $W_1$  and  $W_2$  occur, then the nitrate esters meet the second criterion for the applicability of RRKM theory. However, if the state  $W_3$  intersects the state  $W_1$ , as indicated in the figure, then the molecule has a curve-crossing in the reaction channel. In this case, the intersection involves states of different spins, and coupling between such states is said to be "spin forbidden". To the cogniscenti forbidden is a hollow expletive, since many examples of "spin-forbidden" reactions proceeding at measurable rates exist. The curve-crossing shown, if it is real, will have the effect of accelerating the rate of reaction at high-temperature and pressure. In this case, the pressure dependence of the nitrate ester decomposition cannot be determined by RRKM theory.

The question remains, however, does the curve crossing exist? Harris (18) postulated the crossing on the basis of the diffuse nature of the electronic spectra of nitrate esters, an interpretation then

16. I. Oref, "Nonrandomization of Energy in Unimolecular Reaction. The Effects of High Pressures on the Apparent Rate Constant", J. Chem. Phys., 63, 3138 (1975).
17. G.F. Adams and J.R. Ward, "Assessments of Recent Models for Pressure and Volume Effects on REactions of Ballistic Materials", Tech. Rpt. ARBRL-TR-02055, April 1978. (AD #A054640)
18. L.E. Harris, "The Lower Electronic States of Nitrite and Nitrate Ion, Nitromethane, Nitramide, Nitric Acid, and Nitrate Esters", J. Chem. Phys., 58, 5615 (1973).




Figure 2. Schematic Representation of the Variation of Electronic Energy With Bond Length for Several States of a Nitrate Ester.

considered quite reasonable. Recently, however, a theoretical analysis by Pack (19) demonstrated that there are alternative explanations of diffuse structure in electronic spectra. This being the case, it is not clear that such a crossing occurs in the electronic structure of nitrate esters. Experimental (20) and theoretical (21) investigations of this problem are underway in this laboratory, and a resolution of the problem is anticipated.

In any event, such an esoteric perturbation on the decomposition rate is outside the ranges of validity of the published models for the effects of pressure on ballistic properties, since both models were limited to consideration of the effects of pressure on vibrational frequencies. It is concluded, on the basis of the analysis described in this report, that the initial decomposition reaction of the nitrate esters contained in double-base propellants cannot be the rate-determining step for the observed variation of burning-rate with pressure.

19. R.T. Pack, "Simple Theory of Diffuse Vibrational Structure in Continuous UV Spectra of Polyatomic Molecules. I.", J. Chem. Phys., 65, 4765 (1976).
20. R. Beyer, private communication.
21. G.F. Adams, G.D. Purvis and R.J. Bartlett, unpublished calculations.

## REFERENCES

1. M. M. Ibiricu and F.A. Williams, "Mechanisms for the Steady Deflagration of Double-Base Propellants", 12th JANNAF Combustion Meeting, CPIA Publication Number 273, 1976.
2. D. J. Pastine, M.J. Kamlett and S.J. Jacobs, "Volume and Pressure Dependence of Some Kinetic Processes in Explosives", Naval Surface Weapons Center/WOL/TR75-149, 1975.
3. G.D. Mendenhall, D.M. Golden, and S.W. Benson, "The Very-Low-Pressure Pyrolysis (VLPP) of n-Propyl Nitrate, tert-Butyl Nitrate, and Methyl Nitrite. Rate Constants for Some Alkoxy Radical Reactions", *Int. J. Chem. Kinetics*, 7, 725 (1975).
4. W. Forst, Theory of Unimolecular Reactions, Academic Press, New York, (1973).
5. P.J. Robinson and K.A. Holbrook, Unimolecular Reactions, Wiley-Interscience, New York (1972).
6. F.A. Lindeman, Title not available, *Trans. Faraday Soc.*, 17, 598 (1922).
7. C.N. Hinshelwood, "The Theory of Unimolecular Reactions", *Proc. Roy Soc.(A)*, 113, 230 (1927).
8. O.K. Rice and H.C. Ramsperger, "Theories of Unimolecular Reactions at Low Pressures", *J. Amer. Chem. Soc.*, 49, 1617 (1927).
9. O.K. Rice and H.C. Ramsperger, "Theories of Unimolecular Reactions at Low Pressures. II.", *J. Amer. Chem. Soc.*, 50, 617 (1928).
10. L.S. Kassel, Kinetics of Homogeneous Gas Reactions, Chemical Catalog Co., New York, 1932.
11. R.A. Marcus, "Unimolecular Dissociations and Free Radical Recombination Reactions", *J. Chem. Phys.*, 20, 359 (1952).
12. H.S. Johnston, Gas-Phase Reaction Rate Theory, Ronald, New York, 1966.
13. S.W. Benson, Thermochemical Kinetics, Wiley, New York, 1968.
14. H.E. O'Neal and S.W. Benson, "The Biradical Mechanism in Small Ring Compounds", *J. Phys. Chem.*, 72, 1866, 1968.
15. W.L. Hase and D. Bunker, RRKM, Program No. 234, Quantum Chemistry Program Exchange, University of Indiana, Bloomington, Indiana.

16. I. Oref, "Nonrandomization of Energy in Unimolecular Reaction. The Effects of High Pressures on the Apparent Rate Constant", J. Chem. Phys., 63, 3138 (1975).
17. G.F. Adams and J.R. Ward, "Assessments of Recent Models for Pressure and Volume Effects on Reactions of Ballistics Material", Technical Report ARBRL-TR2055, April 1978. (AD #A054640)
18. L.E. Harris, "The Lower Electronic States of Nitrite and Nitrate Ion, Nitromethane, Nitramide, Nitric Acid and Nitrate Esters", J. Chem. Phys., 58, 5615 (1973).
19. R. T. Pack, "Simple Theory of Diffuse Vibrational Structure in Continuous UV Spectra of Polyatomic Molecules. I.", J. Chem. Phys. 65, 4765 (1976).
20. R. Beyer, private communication.
21. G.F. Adams, G.D. Purvis and R.J. Bartlett, unpublished calculations.

# APPENDIX A. COMMENTS ON DETAILS OF THE RRKM CALCULATION

The working expressions for  $k_{\text{uni}}$  and  $k_a(E^*)$  are repeated here for convenience.

$$k_{\text{uni}} = \frac{L^\ddagger Q_1^\ddagger}{h Q_1 Q_2} \exp(-E_0/RT) \Delta E^+ \sum_{i=1}^{i_{\text{max}}} \left\{ \frac{\{\Sigma P(E_{\text{VR}}^+)\} \exp(-E^+/RT)}{1 + k_a(E^*)/\lambda ZP} \right\}_i \quad (\text{A.1})$$

$$k_a(E^*) = \frac{L^\ddagger Q_1^\ddagger}{h Q_1 N^*(E^*)} \sum_{E_{\text{VR}}^+=0}^{E_{\text{VR}}^+} P(E_{\text{VR}}^+) \quad (\text{A.2})$$

For the sake of completeness, the methods used to evaluate the sums and densities of molecular quantum states should be outlined. Application of the RRKM method is critically dependent in the methods used to evaluate the density of quantum states  $N^*(E_{\text{VR}}^*)$  for the reactant A and the sum of quantum states  $\Sigma P(E_{\text{VR}}^+)$  for the postulated activated complex  $A^\ddagger$ . The computation of  $N^*$  and  $\Sigma P^\ddagger$  for n-propyl nitrate was performed using the semiclassical method developed by Whitten and Rabinovitch (1), and this method will be described briefly.

The density of quantum states  $N(E)$  can be obtained from the sum of quantum states  $W(E) = \Sigma P(E)$  by differentiation with respect to energy. That is

$$N(E) = \frac{dW(E)}{dE} = \frac{d(\Sigma P(E))}{dE}, \quad (\text{A.3})$$

so we need only find a differentiable form for  $W(E)$ . For a set of  $s$  oscillators, each set of quantum numbers  $v_i$  gives rise to one quantum state. The total number of states at all energies up to and including  $E_v$  is given by,

# We shall hereafter adapt the notation  $W(E) = \Sigma P(E)$ , and omit superscripts.

1. G.Z. Whitten and B.S. Rabinovitch, "Accurate and Facile Approximation for Vibrational Energy-Level Sums", J.Chem. Phys. 38, 2466 (1963).

$$\sum_{E_V=0}^{E_V} P(E_V) = \sum_{v_s=0} \sum_{v_{s-1}=0} \dots \sum_{v_1=0} 1 \quad (A.4)$$

where the summation is taken over all  $v_i$  such that the total vibrational energy is less than or equal to  $E_V$ . In the classical approximation, this sum is replaced by an integral over the occupation number,  $v_i$ . The evaluation of the integral is given by Robinson and Holbrook<sup>1(2)</sup>, in Appendix 6, and the result is

$$W(E_V) = P(E_V) \approx \frac{E_V^s}{s! \pi \hbar \nu_i} \quad (A.6)$$

Marcus and Rice (3) extended the classical treatment by suggesting that an oscillator with non-fixed energy  $E_V$  has a total vibrational energy above the ground state of  $E_V + E_z$ , where  $E_z$  is the zero-point vibrational energy. Modification of the classical expressions (A.5) and (2a) yields

$$W(E_V) = \frac{(E_V + E_z)^s}{s! \pi \hbar \nu_i} \quad (A.7)$$

$$N(E_V) = \frac{(E_V + E_z)^{s-1}}{(s-1)! \pi \hbar \nu_i} \quad (A.8)$$

These expressions, though more accurate than the classical approximations, always overestimate the number of states at any level. An improved version of the semiclassical expressions has been developed by Rabinovitch and co-workers.

It was suggested that only a fraction of the zero-point energy should be added to the vibrational energy  $E_V$ . An empirical factor  $a$  was introduced to multiply  $E_z$  in equations (A.7) and (A.8). Whitten and Rabinovitch discovered a relation between the parameter,  $a$ , and a modified frequency dispersion parameter,  $\beta$ , defined by

$$\beta = \frac{s-1}{s} \frac{\langle v^2 \rangle}{\langle v \rangle^2} \quad (A.9)$$

The relation is

$$a = 1 - \beta \cdot \omega(E) \quad (A.10)$$

2. P.J. Robinson and K.A. Holbrook, Unimolecular Reactions, Wiley-Interscience, New York (1972).
3. R.A. Marcus and O.K. Rice, "The Kinetics of the Recombination of Methyl Radicals and Iodine Atoms", J. Phys. and Colloid. Chem., 55, 894 (1951).



where  $E' = E_v / E_z$ , and  $\omega(E')$  is a function of  $E'$  defined by

$$(0.1 < E' < 1.0) \quad \omega = (5.E' + 2.73 E'^{0.5} + 3.51)^{-1} \quad (\text{A.11a})$$

$$(1.0 < E' < 8.0) \quad \omega = \exp(-2.4191 E'^{0.25}) \quad (\text{A.11b})$$

The molecular frequencies are used to calculate  $\beta$  and  $E_z$ , and then for a non-fixed energy  $E_v$ , the reduced energy  $E'$  is calculated and used to obtain  $\omega(E')$ . Given this data, the parameter  $a$  and the sum of states can be computed. The density of states can then be computed using the expression

$$N(E_v) = \frac{(E_v + aE_z)^{s-1}}{(s-1)! \pi h \nu_i} \left[ 1 - \beta \left( \frac{d\omega}{dE'} \right) \right]$$

where  $\left( \frac{d\omega}{dE'} \right)$  is obtained by differentiating (A.11a) and (A.11b).

The Whitten Rabinovitch method has been thoroughly tested, and the results are in good agreement with the direct-count for non-fixed energies  $E_v$  greater than  $E_z$ , and often for lower energies. Whitten and Rabinovitch have extended the treatment to include the case where active rotations need to be taken into account. Again, the equations suggested have been widely tested and the results appear to be good approximations to  $N(E_{vr})$  and  $N(E_{vr})$ .

#### APPENDIX B: DATA FORMAT FOR RRKM

The computer code RRKM is stored as an UPDATE file on the BRL Cyber 173. This code can be used by submitting the following deck of cards.

1. JOBCARD
2. ACCOUNTING CARD
3. ATTACH (OLDPL,RRKM,ID-GEOFADAMS)
4. UPDATE(F)
5. FTN,I.
6. LGO.
7. \*EOR
8. \* ID,(MARY) ← arbitrary choice of ID
9. \*EOR
- DATA
10. \*EOI.

This job will produce values for  $N(E^*)$ ,  $w(E^*)$  and  $k_a(E^*)$ . These data

can be used to calculate  $k_{\text{uni}}$  as a function of pressure by solving equation (A.1). Unless one has substantial experience with such computations, it is recommended that the computation of  $k_{\text{uni}}$  be carried out on a desk-top computer or as an interactive exercise on a terminal. By following this procedure, one need not compute  $k_{\text{uni}}$  for an excessive pressure range. Users are, however, invited to disagree with this philosophy.

The data required to perform a standard RRKM calculation is as follows:

1. Format 3F 10.0, 15, F10.0. A card with  $E_0$  (threshold energy),  $E^*$  (lowest molecular energy used in the calculations),  $\Delta E^+$  (upward increment of same),  $n$ , the number of successive molecular energies for which the calculation will be made, and  $\sigma_p$ , the statistical factor.
2. Two title cards. Say anything you like.
3. Format 2I10.  $n_f$ , the number of reactant frequencies and  $n_r$ , the number of free internal rotors.
4. Format 7F10.0. The  $n_f$  molecular frequencies, seven per card. Units,  $\text{cm}^{-1}$ .
5. Reduced moments of inertia for the internal rotors of the reactant. Units  $\text{amu } \text{\AA}^2$ . Format 7F10.0. Omit this card if  $N_r = 0$ .
6. Symmetry numbers corresponding to item 5. Same format. Omit if  $n_\sigma = 0$ .
7. Format, I10. KROT=0.
8. FORMAT, 3F.10. Three overall moments of inertia for the reactant; units  $\text{amu-}\text{\AA}^2$ .
9. FORMAT, I10. NANH=0.
10. FORMAT, 2I10. JDEN=0, JSUM=0.
11. FORMAT, I10. NCRIT=0.
12. FORMAT, 2I10.  $n_f^+$  and  $n_r^+$ , same as item 3 but for the critical configuration.  $n_f^+ + n_r^+ = N_f + N_r - 1$ .
13. Frequencies, like item 4 except for critical configuration.

14. Reduced moments, like item 5, except for critical configuration.

15. Symmetry numbers, like item 6, except for critical configuration.

16. The overall moments of inertia for the critical configuration as in item 8.  $I_x$ ,  $I_y$ , and  $I_z$  sequence must be the same as for the molecule.

17. Format F10.0. Rotational temperature for adiabatic rotations, <sup>o</sup>K.

18. Format I10. NTRAN = 0.

Input may be stacked. To stop the program add a blank card after item 18.

## REFERENCES

1. G.Z. Whitten and B.S. Rabinovitch, "Accurate and Facile Approximation for Vibrational Energy-Level Sums", J. Chem. Phys. 38, 2466 (1963).
2. P.J. Robinson and K.A. Holbrook, Unimolecular Reactions, Wiley-Interscience, New York (1972).
3. R.A. Marcus and O.K. Rice, "The Kinetics of the Recombination of Methyl Radicals and Iodine Atoms", J. Phys. and Colloid. Chem., 55, 894 (1951).

# DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
12	Commander Defense Documentation Center ATTN: DCC-TCA Cameron Station Alexandria, VA 22314	1	Commander US Army Communications Rsch and Development Command ATTN: DRDCO-SGS Ft. Monmouth, NJ 07703
1	Director Defense Advanced Research Projects Agency ATTN: C.R. Lehner 1400 Wilson Boulevard Arlington, VA 22209	1	Commander US Army Missile Research and Development Command ATTN: DRDMI-R Redstone Arsenal, AL 35809
2	Director Institute for Defense Analyses ATTN: H. Wolfhard R.T. Oliver 400 Army-Navy Drive Arlington, VA 22202	1	Commander US Army Missile Materiel Readiness Command ATTN: DRSMI-AOM Redstone Arsenal, AL 35809
1	Commander US Army Materiel Development and Readiness Command ATTN: DRCDMD-ST, N. Klein 5001 Eisenhower Avenue Alexandria, VA 22333	1	Commander US Army Tank Automotive Research and Development Command ATTN: DRDTA-UL Warren, MI 48090
1	Commander US Army Aviation Research and Development Command ATTN: DRSAB-E P.O. Box 209 St. Louis, MO 63166	1	Commander US Army Armament Materiel Readiness Command ATTN: DRSAR-LEP-L, Tech Lib Rock Island, IL 61299
1	Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035	5	Commander US Army Armament Research & Development Command ATTN: DRDAR-TSS (2 cys) DRDAR-LC, Dr. Kemmy Dr. Jack Alster Dr. J. Sharma Dover, New Jersey 07801
1	Commander US Army Electronics Research & Development Command Technical Support Activity ATTN: DELSD-L Ft. Monmouth, NJ 07703	6	Commander US Army Armament Research and Development Command ATTN: DRDAR-LCE-CI, J. Lannon DRDAR-LC, J.P. Picard DRDAR-LCE, C. Lenchitz DRDAR-LCE, R.F. Walker DRDAR-SCA-CC, C. Dickey DRDAR-SCA-PP, L. Stiefel Dover, New Jersey 07801

# DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
1	Chief, Benet Weapons Lab LCWSL, USAARRADCOM ATTN: DRDAR-LCB-RD, R. Thierry Watervliet, NY 12189	2	Commander US Naval Surface Weapons Center ATTN: S.J. Jacobs/Code 240 Code 730 Silver Spring, MD 20910
1	Commander US Army White Sands Missile Range ATTN: STEWS-VT White Sands, NM 88002	1	Commander US Naval Surface Weapons Center ATTN: Library Br, DX-21 Dahlgren, VA 22448
1	Commander US Army Materials and Mechanics Research Center ATTN: DRXMR-ATL Watertown, MA 02172	1	Commander US Naval Underwater Systems Center Energy Conversion Department ATTN: R.S. Lazar/Code 5B331 Newport, RI 02840
1	Commander US Army Natick Research and Development Command ATTN: DRXRE, D. Sieling Natick, MA 01762	2	Commander US Naval Weapons Center ATTN: R. Derr C. Thelen China Lake, CA 93555
1	Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SL, Tech Lib WSMR, NM 88002	1	Commander US Naval Research Laboratory ATTN: Code 6180 Washington, DC 20375
1	Commander US Army Research Office ATTN: Tech Lib P.O. Box 12211 Research Triangle Park, NC 27706	3	Superintendent US Naval Postgraduate School ATTN: Tech Lib David Netzer Allen Fuhs Monterey, CA 93940
1	Chief of Naval Research ATTN: Code 473 800 N. Quincy Street Arlington, VA 22217	2	Commander US Naval Ordnance Station ATTN: A. Roberts Tech Lib Indian Head, MD 20640
1	Commander US Naval Sea Systems Command ATTN: J.W. Murrin (NAVSEA-0331) National Center, Bldg. 2, Rm 6E08 Washington, DC 20360	2	AFOSR ATTN: J.F. Masi B.T. Wolfson Bolling AFB, DC 20332

# DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
3	AFRPL (DYSC/D. George, J.N. Levine, W. Roe) Edwards AFB, CA 93523	1	General Electric Company Armament Department ATTN: M. J. Bulman Lakeside Avenue Burlington, VT 05402
1	Jet Propulsion Laboratory ATTN: Leon Stromd, Bldg 125 4800 Oak Grove Drive Pasadena, CA 91103	2	Hercules Incorporated Alleghany Ballistic Laboratory ATTN: R. Miller Tech Lib Cumberland, MD 21501
1	Aerojet Solid Propulsion Co. ATTN: P. Micheli Sacramento, CA 95813	1	Hercules Incorporated Bacchus Works ATTN: M. Beckstead Magna, UT 84044
1	ARO Incorporated ATTN: N. Dougherty Arnold AFS, TN 37389	1	Hercules Incorporated Eglin AFB Laboratory ATTN: R. L. Simmons P. O. Box 1646 Eglin AFB, FL 32542
1	Atlantic Research Corporation ATTN: M. K. King 5390 Cherokee Avenue Alexandria, VA 22314	1	IITRI ATTN: M. J. Klein 10 West 35th Street Chicago, IL 60615
1	AVCO Corporation AVCO Everett Research Lab Div ATTN: D. Stickler 2385 Revere Beach Parkway Everett, MA 02149	1	Lockheed Palo Alto Rsch Labs ATTN: Tech Info Ctr 3521 Hanover Street Palo Alto, CA 94304
2	Calspan Corporation ATTN: E. B. Fisher A. P. Trippe P.O. Box 235 Buffalo, NY 14221	1	Olin Corporation Badger Army Ammunition Plant ATTN: J. Ramnarace Baraboo, WI 53913
1	ENKI Corporation ATTN: M. I. Madison 9015 Fulbright Avenue Chatsworth, CA 91311	1	Olin Corporation New Haven Plant ATTN: R. L. Cook D. W. Riefler 275 Winchester Avenue New Haven, CT 06504
1	Foster Miller Associates, Inc. ATTN: A. J. Erickson 135 Second Avenue Waltham, MA 02154	1	Paul Gough Associates, Inc. ATTN: P. S. Gough P. O. Box 1614 Portsmouth, NH 03801
1	General Electric Flight Propulsion ATTN: Tech Lib Cincinnati, OH 45215		

# DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
1	Physics International Company 2700 Merced Street Leandro, CA 94577	2	Thiokol Corporation Wasatch Division ATTN: John Peterson Tech Lib P.O. Box 524 Brigham City, UT 84302
1	Pulsepower Systems, Inc. ATTN: L.C. Elmore 815 American Street San Carlos, CA 94070	1	TRW Systems Group ATTN: H. Korman One Space Park Redondo Beach, CA 90278
1	Science Applications, Inc. ATTN: R.B. Edelman 23146 Cumorah Crest Woodland Hills, CA 91364	2	United Technology Center ATTN: R. Brown Tech Lib P.O. Box 358 Sunnyvale, CA 94088
2	Rockwell International Corp. Rocketdyne Division ATTN: C. Obert J.E. Flanagan 6633 Canoga Avenue Canoga Park, CA 91304	1	Universal Propulsion Co. ATTN: H.J. McSpadden P.O. Box 546 Riverside, CA 92502
2	Rockwell International Corp. Rocketdyne Division ATTN: W. Haymes Tech Lib McGregor, TX 76657	1	Battelle Memorial Institute ATTN: Tech Lib 505 King Avenue Columbus, OH 43201
1	Shock Hydrodynamics, Inc. ATTN: W.H. Anderson 4710-16 Vineland Avenue North Hollywood, CA 91602	1	Brigham Young University Dept of Chemical Engineering ATTN: R. Coates Provo, UT 84601
1	Thiokol Corporation Elkton Division ATTN: E. Sutton Elkton, MD 21921	1	California Institute of Tech 204 Karman Lab Mail Stop 301-46 ATTN: F.E.C. Culick 1201 E. California Street Pasadena, CA 91125
3	Thiokol Corporation Huntsville Division ATTN: D. Flanigan R. Glick Tech Lib Huntsville, AL 35807	1	Case Western Reserve Univ. Division of Aerospace Sciences ATTN: J. Tien Cleveland, OH 44135



# DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
3	Georgia Institute of Technology School of Aerospace Engineering ATTN: B. T. Zinn E. Price W. C. Strahle Atlanta, GA 30332	2	Purdue University School of Mechanical Engineering ATTN: J. Osborn S. N. B. Murthy TSPC Chaffee Hall West Lafayette, IN 47906
1	The Johns Hopkins University Applied Physics Laboratory Chemical Propulsion Information Agency ATTN: T. Christian Johns Hopkins Road Laurel, MD 20810	1	Southwest Research Institute Fire Research Section ATTN: W. H. McLain P.O. Drawer 28510 San Antonio, TX 78228
1	Lehigh University Department of Mechanical Engineering ATTN: A. MacPherson Allentown, PA 18015	1	Stanford Research Institute Propulsion Sciences Division ATTN: Tech Lib 333 Ravenswood Avenue Menlo Park, CA 94024
1	Massachusetts Institute of Technology Department of Mechanical Engineering ATTN: T. Toong Cambridge, MA 02139	1	Stevens Institute of Technology Davidson Laboratory ATTN: R. McAlevy, III Hoboken, NJ 07030
1	Pennsylvania State University Applied Research Laboratory ATTN: G. M. Faeth P.O. Box 30 State College, PA 16801	1	Texas Technological College Department of Chemistry ATTN: Dr. T. O'Brien Lubbock, TX 79406
1	Pennsylvania State University Department of Mechanical Engineering ATTN: K. Kuo University Park, PA 16801	1	University of California, San Diego AMES Department ATTN: F. Williams P.O. Box 109 La Jolla, CA 92037
2	Princeton University Forrestal Campus Library ATTN: L. Caveny Tech Lib P.O. Box 710 Princeton, NJ 08540	1	University of Illinois Department of Aeronautical Engineering ATTN: H. Krier Transportation Building, Rm 105 Urbana, IL 61801

DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>
1	University of Minnesota Dept of Mechanical Engineering ATTN: E. Fletcher Minneapolis, MN 55455
2	University of Utah Dept of Chemical Engineering ATTN: A. Baer G. Flandro Salt Lake City, UT 84112

Aberdeen Proving Ground

Dir, USAMSAA

Cdr, USATECOM

ATTN: DRSTE-SG-H